

Commutative Rings of Differential Operators Connected with Two-Dimensional Abelian Varieties

A.E. Mironov *

Abstract

In this paper we find the explicit formulas of two dimensional commuting (2×2) -matrix differential operators which were introduced by Nakayashiki. The common eigen functions and eigen values of these operators are parametrized by the points of principally polarized Abelian varieties.

1 Introduction

The main result of this paper is construction of two-dimensional (2×2) -matrix differential operators with doubly periodic coefficients from spectral functions and the vector Baker–Akhiezer eigenfunction that was introduced by Nakayashiki in [1]. A remarkable property of these operators is the fact that they are finite-gap at every energy level. Namely, their Blöch functions, the eigenfunctions of both the differential operator and the operators of translation by periods, are parametrized by the points of a two-dimensional principally polarized Abelian variety X^2 . Moreover, the eigenvalues (making the spectrum) are given by some meromorphic function $\lambda(z)$ on X^2 with pole on a theta divisor. We indicate a procedure for constructing a commutative ring of such operators from an Abelian variety X^2 (with an irreducible theta divisor) and a spectral function λ .

The multidimensional inverse problem was first solved by B. A. Dubrovin, I. M. Krichever, and S. P. Novikov for a periodic Schrödinger operator with

*Institute of Mathematics, 630090 Russia, Novosibirsk; e-mail: mironov@math.nsc.ru

a magnetic field [2]. They described a procedure for constructing such operators that are finite-gap at one energy level, i.e., their Bloch functions with a fixed eigenvalue E_0 are parametrized by the points of a Riemann surface of finite genus. Later, A. P. Veselov and S. P. Novikov [3] distinguished potential operators (i.e., operators with zero magnetic field) among them.

The finite-gap property of the Schrödinger operator at a fixed energy level is an exceptional phenomenon: Feldman, Knorrer, and Trubowitz [4] demonstrated that the two-dimensional Schrödinger operator with a smooth real potential can be finite-gap only at one energy level. As was first observed by Sato, for matrix operators it is possible that the Bloch functions are parametrized by surfaces of finite genus (in our case by the level surfaces $\lambda = \text{const}$) for all energy levels. This idea was implemented by Nakayashiki [1, 5].

In [1], using the Fourier–Mukai transform method [6], Nakayashiki constructed the Baker–Akhiezer module over a ring of differential operators in g space variables, where g is the dimension of a principally polarized Abelian variety X^g with a nonsingular theta divisor. To this module there corresponds (up to conjugation) a commutative ring of $(g! \times g!)$ -matrix partial differential operators whose coefficients are, in general, defined locally. Each operator corresponds to some meromorphic function on X^g with pole on the theta divisor (a spectral function) which parametrizes the eigenvalues of the operator (these operators are now referred to as the Nakayashiki operators). The vector-function whose components are the elements of the basis for the Baker–Akhiezer module parametrizes the general eigenfunctions of these operators.

Observe that this construction requires nonsingularity of the theta divisor. As was demonstrated by Andreotti and Mayer [7], a theta divisor of a general Abelian variety is a nonsingular subvariety; however, a theta divisor of the Jacobian varieties of Riemann surfaces has singularities for $g > 3$ and in the case of hyperelliptic surfaces also for $g = 3$.

In Section 2 we describe the Fourier–Mukai transform [6] in the needed situation, recall Nakayashiki’s construction of the Baker–Akhiezer module [1], and indicate connection between Krichever’s construction [8] and the Fourier–Mukai transform [6].

In Section 3 we give a new short analytical proof of Nakayashiki’s theorem of freeness of the Baker–Akhiezer module in dimension 2. It essentially uses the fact that the Abelian variety X^2 in question is two-dimensional unlike the proof of the general case in [1] and requires a minimal apparatus of algebraic geometry. Here we also introduce a basis for the Baker–Akhiezer module

for $g = 2$ in which we manage to find the coefficients of the Nakayashiki operators.

In Section 4 we describe an effective procedure for constructing the Nakayashiki operators for $g = 2$. In Propositions 1–3 we obtain explicit formulas for operators generating, as proven in Lemma 9, the whole ring of the Nakayashiki operators. In Proposition 1 we introduce commuting operators $Z_1, \dots, Z_g$ such that, for every Nakayashiki operator L , the commutator $[L, Z_j]$ is a Nakayashiki operator too. In Proposition 2 we find explicit formulas for the second-order Nakayashiki operators and in Proposition 3, explicit formulas for the operators Z_j .

The author is grateful to S. P. Novikov and I. A. Taimanov for posing the problem and to I. A. Taimanov for useful discussions and remarks.

2 Nakayashiki's Construction

Let $X^g = \mathbf{C}^g / (\mathbf{Z}^g + \Omega \mathbf{Z}^g)$ be a principally polarized Abelian complex variety, where Ω is a symmetric $(g \times g)$ -matrix with $\text{Im} \Omega > 0$. Denote by $\text{Pic}^0(X^g)$ the Picard variety of X^g . If X^g is principally polarized then $\text{Pic}^0(X^g)$ is isomorphic to X^g . The isomorphism is given by the mapping $z \rightarrow x$, where $z \in X^g$ and $x \in \text{Pic}^0(X^g)$. The sections of a bundle x , lifted to the universal covering $\mathbf{C}^g$, are given by functions $h(z)$ satisfying the periodicity conditions

$$h(z + \Omega m + n) = \exp(-2\pi i \langle m, x \rangle) h(z),$$

where $m, n \in \mathbf{Z}^g$ and $\langle m, x \rangle = m_1 x_1 + \dots + m_g x_g$.

Denote by $\mathcal{P}$ the Poincaré bundle over $X^g \times \text{Pic}^0(X^g)$. It is determined by the following properties. The bundle corresponding to $x \in \text{Pic}^0(X^g)$ is isomorphic to $\mathcal{P}|_{X^g \times \{x\}}$ and the bundles $\mathcal{P}|_{X^g \times \{0\}}$ and $\mathcal{P}|_{\{0\} \times \text{Pic}^0(X^g)}$ are trivial. The sections of $\mathcal{P}$, lifted to the universal covering $\mathbf{C}^g \times \mathbf{C}^g$, are given by functions $f(z, x)$ such that

$$f(z + \Omega m_1 + n_1, x + \Omega m_2 + n_2) = \exp(-2\pi i (\langle m_1, x \rangle + \langle m_2, z \rangle)) f(z, x), \quad (1)$$

where $m_j, n_j \in \mathbf{Z}^g$. Denote by $z = (z_1, \dots, z_g)^\top \in \mathbf{C}^g$ the coordinates of points on the universal covering of X^g and by $x = (x_1, \dots, x_g)^\top \in \mathbf{C}^g$, the coordinates of points on the universal covering of $\text{Pic}^0(X^g)$. We identify the sections of $\mathcal{P}$ with the functions on $\mathbf{C}^g \times \mathbf{C}^g$ satisfying (1).

The theta function with characteristic $[a, b]$ is defined by the series

$$\theta[a, b](z, \Omega) = \sum_{n \in \mathbf{Z}^g} \exp(\pi i \langle \Omega(n + a), (n + a) \rangle + 2\pi i \langle (n + a), (z + b) \rangle),$$

where $a, b \in \mathbf{C}^g$. For short, the function $\theta(z) = \theta[0, 0](z, \Omega)$ is referred to as a *theta function*. The function $\theta[a, b](z, \Omega)$ possesses the periodicity properties

$$\theta[a, b](z + m, \Omega) = \exp(2\pi i \langle a, m \rangle) \theta[a, b](z, \Omega),$$

$$\theta[a, b](z + \Omega m, \Omega) = \exp(-2\pi i \langle b, m \rangle - \pi i \langle m \Omega, m \rangle - 2\pi i \langle m, z \rangle) \theta[a, b](z, \Omega),$$

$m \in \mathbf{Z}^g$.

Denote by Θ the zero set of a theta function (a theta divisor) which is a subvariety in X^g .

Given a subvariety $Y \subset X^g$ of codimension 1, we let A_Y stand for the space of meromorphic functions on X^g with pole in Y . We denote by $F_Y(U)$ the space of meromorphic sections of $\mathcal{P}$ over subsets of the form $X^g \times U \subset X^g \times \text{Pic}^0(X^g)$ which have pole in $Y \times U$, where U is an open subset in $\text{Pic}^0(X^g)$.

The set $F_Y = \bigcup_U F_Y(U)$ is called the Fourier–Mukai transform of A_Y [6].

In [1] Nakayashiki constructed “covariant differentiation” operators (a connection on $\mathcal{P}$)

$$\nabla_j : F_Y(U) \rightarrow F_Y(U), \quad \nabla_k \nabla_j = \nabla_j \nabla_k, \quad k, j = 1, \dots, g,$$

that give $F_Y(U)$ a module structure over the ring $\mathcal{O}_U[\nabla_1, \dots, \nabla_g]$, where $U \subset \text{Pic}^0(X^g)$ is an open subset and $\mathcal{O}_U$ is the ring of analytic functions on U . By construction, $F_Y(U)$ is also an A_Y -module.

Nakayashiki introduced the Baker–Akhiezer functions

$$f(z, x) \exp\left(-\sum_{j=1}^g x_j \partial_{z_j} \log \theta(z)\right), \quad f(z, x) \in F_\Theta.$$

The operators ∇_j are defined by the formula

$$\nabla_j = \partial_{x_j} - \partial_{z_j} \log \theta(z).$$

Let M_c denote the space of the Baker–Akhiezer functions

$$\left\{ f(z, c + x) \exp\left(-\sum_{j=1}^g x_j \partial_{z_j} \log \theta(z)\right) \mid f(z, c + x) \in \bigcup_U F_\Theta(c + U) \right\},$$

where $U \subset \mathbf{C}^g$ is a neighborhood of zero, $c \in \mathbf{C}^g$, and $f(z, c+x)$ belongs to the range of the Fourier–Mukai transform F_Θ . Each function $f(z, c+x)$ is defined on $\mathbf{C}^g \times U$ and has pole only in $\Theta \times U$. Roughly speaking, $f(z, c+x)$ is a germ of a function at $x=0$ with respect to x . It follows from (1) that

$$f(z + \Omega m + n, c+x) = \exp(-2\pi i \langle m, (c+x) \rangle) f(z, c+x).$$

We denote by $M_c(k)$ the subset of functions in M_c such that $f(z, c+x)$ has a pole of order $\leq k$ in $\Theta \times U$. The definition implies that M_c is a module over the ring $\mathcal{D} = \mathcal{O}[\partial_{x_1}, \dots, \partial_{x_g}]$ of differential operators (a $\mathcal{D}$ -module), where $\mathcal{O}$ is the ring of analytic functions in $x_1, \dots, x_g$ in a neighborhood of $0 \in \mathbf{C}^g$. The $\mathcal{D}$ -module M_c is referred to as the *Baker–Akhiezer module*. From the definition of the Fourier–Mukai transform we infer that M_c also possesses the structure of an A_Θ -module. The $\mathcal{D}$ -module M_c can be described by means of theta functions [1].

Lemma 1. *The equality holds:*

$$M_c = \sum_{n=1}^{\infty} \sum_{a \in \mathbf{Z}^g / n\mathbf{Z}^g} \mathcal{O} \cdot \frac{\theta[\frac{a}{n}, 0](nz + c + x, n\Omega)}{\theta^n(z)} \exp\left(-\sum_{k=1}^g x_k \partial_{z_k} \log \theta(z)\right).$$

We also need the following [1]:

Lemma 2. *The identity holds:*

$$\begin{aligned} & \partial_{x_j} \left(\frac{\theta[\frac{a}{n}, 0](nz + c + x, n\Omega)}{\theta^n(z)} \exp\left(-\sum_{k=1}^g x_k \partial_{z_k} \log \theta(z)\right) \right) \\ &= \frac{1}{n} \partial_{z_j} \left(\frac{\theta[\frac{a}{n}, 0](nz + c + x, n\Omega)}{\theta^n(z)} \right) \exp\left(-\sum_{k=1}^g x_k \partial_{z_k} \log \theta(z)\right), \end{aligned}$$

$j = 1, \dots, g$.

The next assertion was proven in [1].

Nakayashiki’s Theorem. *If Θ is a nonsingular variety and $c \neq 0$ then M_c is a free $\mathcal{D}$ -module of rank $g!$.*

Fix a basis $\Phi_c = (\phi_{1c}(z, x), \dots, \phi_{g!c}(z, x))^T$ for the $\mathcal{D}$ -module M_c . Suppose that $\lambda(z) \in A_\Theta$. Since M_c is a free $\mathcal{D}$ -module, there exist unique operators $[L_{\Phi_c}(\lambda)]_{kj} \in \mathcal{D}$ such that

$$\sum_{j=1}^{g!} [L_{\Phi_c}(\lambda)]_{kj} \phi_{jc} = \lambda \phi_{kc}, \quad k = 1, \dots, g!.$$

Consequently,

$$L_{\Phi_c}(\lambda)\Phi_c = \lambda\Phi_c, \quad (2)$$

with $[L_{\Phi_c}(\lambda)]_{kj}$ the entries of the matrix differential operator $L_{\Phi_c}(\lambda)$ and $\lambda\Phi_c = (\lambda\phi_{1c}, \dots, \lambda\phi_{g!c})^\top$. Since $L_{\Phi_c}(\lambda)$ are differential operators in x_j while λ depends only on z , from (2) we obtain the commutation condition

$$L_{\Phi_c}(\lambda\mu) = L_{\Phi_c}(\lambda)L_{\Phi_c}(\mu) = L_{\Phi_c}(\mu)L_{\Phi_c}(\lambda),$$

where $\mu(z) \in A_\Theta$. We arrive at the following

Corollary [1]. *There is a ring embedding*

$$L_{\Phi_c} : A_\Theta \rightarrow \text{Mat}(g!, \mathcal{D}),$$

where $\text{Mat}(g!, \mathcal{D})$ is the ring of $(g! \times g!)$ -matrix differential operators. The image of the embedding is a commutative ring of differential operators.

As indicated in [5], Nakayashiki's construction generalizes Krichever's construction [8] which can be interpreted in terms of the Fourier–Mukai transform as follows:

Recall the construction of the Baker–Akhiezer function [8]. Let Γ be a Riemann surface of genus g , let $D = p_1 + \dots + p_g$ be a nonspecial positive divisor on Γ , and let ∞ be a point on Γ other than the points of the divisor. Take a local parameter k^{-1} at ∞ so that $k^{-1}(\infty) = 0$. The *one-point Baker–Akhiezer function with spectral data* $\{\Gamma, \infty, p_1, \dots, p_g, k^{-1}\}$ is a function $\psi(z, x)$, $z \in \Gamma$, defined to within a factor depending only on x , by the following properties:

- (1) $\psi(z, x)$ is meromorphic on $\Gamma \setminus \infty$ and the poles of ψ are independent of x and coincide with $\{p_1, \dots, p_g\}$;
- (2) the function $\psi(z, x) \exp(-kx)$ is analytic in a neighborhood of ∞ .

This function has the form

$$\psi(z, x) = \frac{\theta\left(\mathcal{A}(z) - \sum_{j=1}^g \mathcal{A}(p_j) - \Delta + Vx\right)}{\theta\left(\mathcal{A}(z) - \sum_{j=1}^g \mathcal{A}(p_j) - \Delta\right)} \exp\left(2\pi i x \int_{p_0}^z \eta\right),$$

where $\mathcal{A} : \Gamma \rightarrow X^g$ is the Abel mapping with a basepoint p_0 , X^g is the Jacobian variety of Γ , Δ is the vector of Riemann constants, and $V \in \mathbf{C}^g$ is a vector determined from the spectral data.

The function

$$\frac{\theta\left(\mathcal{A}(z) - \sum_{j=1}^g \mathcal{A}(p_j) - \Delta + Vx\right)}{\theta\left(\mathcal{A}(z) - \sum_{j=1}^g \mathcal{A}(p_j) - \Delta\right)}$$

is the restriction to $\mathcal{A}(\Gamma) \times \{Vx\}$ of the function

$$\frac{\theta\left(\tilde{z} - \sum_{j=1}^g \mathcal{A}(p_j) - \Delta + \tilde{x}\right)}{\theta\left(\tilde{z} - \sum_{j=1}^g \mathcal{A}(p_j) - \Delta\right)}$$

which belongs to the range of the Fourier–Mukai transform $F_{\Theta'}$, where $\Theta' \subset X^g$ is the subvariety defined by the equation

$$\theta\left(\tilde{z} - \sum_{j=1}^g \mathcal{A}(p_j) - \Delta\right) = 0.$$

Denote by $\tilde{\mathcal{D}}(\psi(z, x))$ the $\tilde{\mathcal{D}}$ -module $\{d\psi(z, x) \mid d \in \tilde{\mathcal{D}}\}$, where $\tilde{\mathcal{D}}$ is the ring of differential operators in x . As shown in [8], $\tilde{\mathcal{D}}(\psi(z, x))$ is a free $\tilde{\mathcal{D}}$ -module; moreover, for every meromorphic function $f(z)$ on Γ with a sole pole at ∞ , there is a unique differential operator $L(f)$ such that

$$L(f)\psi(z, x) = f(z)\psi(z, x).$$

Hence, we obtain a correspondence between the spectral data of commutative rings of scalar differential operators in the Burchnell–Chaundy–Krichever theory and the spectral data of commutative rings of the Nakayashiki matrix differential operators:

$$\{\Gamma, \infty, D, f\} \longleftrightarrow \{X^g, \Theta, c, \lambda\}.$$

3 Proof of Nakayashiki’s Theorem for $g = 2$

Here we give a new proof of Nakayashiki’s theorem for $g = 2$. Introduce the following functions in M_c :

$$\psi = \frac{\theta(z + c + x)}{\theta(z)} \exp(-x_1 \partial_{z_1} \log \theta(z) - x_2 \partial_{z_2} \log \theta(z)),$$

$$\psi_{c'} = \frac{\theta(z+c+c'+x)\theta(z-c')}{\theta^2(z)} \exp(-x_1 \partial_{z_1} \log \theta(z) - x_2 \partial_{z_2} \log \theta(z)),$$

and define the surface

$$\Gamma_{c'} = \{z \in \mathbf{C}^2 \mid \theta(z-c') = 0\}.$$

In Lemmas 4 and 5 we prove that $\mathcal{D}(\psi, \psi_{c'}) = \{d_1\psi + d_2\psi_{c'} \mid d_1, d_2 \in \mathcal{D}\}$ is a free $\mathcal{D}$ -module. In Lemma 6 we show that the $\mathcal{D}$ -modules $\mathcal{D}(\psi, \psi_{c'})$ and M_c coincide.

Henceforth subscripts of a theta function denote differentiation with respect to the corresponding variable: $\theta_j(z) = \partial_{z_j} \theta(z)$ and $\theta_{kj}(z) = \partial_{z_k} \partial_{z_j} \theta(z)$.

Lemma 3. *The functions $a\theta_1(z) + b\theta_2(z)$ and $\theta(z+e)$, where $a, b \in \mathbf{C}$ and $e \in \mathbf{C}^2$, have two zeros on Θ with multiplicity counted to within elements of the lattice $\mathbf{Z}^2 + \Omega\mathbf{Z}^2$.*

PROOF. Let X^2 be the Jacobian variety of some Riemann surface Γ of genus 2. Denote by $\widehat{\Gamma}$ the universal covering of Γ . The fundamental group $\pi_1(\Gamma)$ acts on $\widehat{\Gamma}$ and we denote a fundamental domain by S . Let $\mathcal{A} : \widehat{\Gamma} \rightarrow X^2$ be the Abel mapping. The range $\mathcal{A}(\widehat{\Gamma})$ has the equation $\theta(z-\Delta) = 0$. It is easy to verify that the sought number of zeros equals

$$\frac{1}{2\pi i} \int_{\partial S} d \log(a\theta_1(\mathcal{A}(p)-\Delta) + b\theta_2(\mathcal{A}(p)-\Delta)) = \frac{1}{2\pi i} \int_{\partial S} d \log \theta(\mathcal{A}(p)+e-\Delta) = 2.$$

Lemma 3 is proven.

Lemma 4. *If $c' \in \mathbf{C}^2$ and $c' \notin \mathbf{Z}^2 + \Omega\mathbf{Z}^2$ then $\psi_{c'}$ does not belong to the set of functions*

$$\{\alpha_1(x)\partial_{x_1}\psi + \alpha_2(x)\partial_{x_2}\psi + \alpha_3(x)\psi \mid \alpha_1(x), \alpha_2(x), \alpha_3(x) \in \mathcal{O}\}.$$

PROOF. Suppose that the assertion of the lemma is false. Then there exist $\alpha_1(x), \alpha_2(x), \alpha_3(x) \in \mathcal{O}$ such that

$$\alpha_1(x)\partial_{x_1}\psi + \alpha_2(x)\partial_{x_2}\psi + \alpha_3(x)\psi = \psi_{c'}.$$

Thereby the following equality holds for $z \in \Gamma_{c'}$:

$$\alpha_1(x) \left(\frac{\theta_1(z+c+x)}{\theta(z)} - \frac{\theta(z+c+x)\theta_1(z)}{\theta^2(z)} \right)$$

$$+\alpha_2(x) \left(\frac{\theta_2(z+c+x)}{\theta(z)} - \frac{\theta(z+c+x)\theta_2(z)}{\theta^2(z)} \right) + \alpha_3(x) \frac{\theta(z+c+x)}{\theta(z)} = 0.$$

Let p be an intersection point of $\Gamma_{c'}$ and Θ (Lemma 3). It follows from the last equality that

$$\alpha_1(x)\theta_1(p) + \alpha_2(x)\theta_2(p) = 0.$$

Since Θ is a smooth Riemann surface, we have $\theta_1(p) \neq 0$ or $\theta_2(p) \neq 0$. For definiteness, assume that $\theta_1(p) \neq 0$. Then

$$\frac{\alpha_1(x)}{\alpha_2(x)} = -\frac{\theta_2(p)}{\theta_1(p)}.$$

Consequently,

$$-\frac{\theta_2(p)}{\theta_1(p)} \partial_{x_1} \log \theta(z+c+x) + \partial_{x_2} \log \theta(z+c+x) = -\frac{\theta_2(p)}{\theta_1(p)} \frac{\theta_1(z)}{\theta(z)} + \frac{\theta_2(z)}{\theta(z)} - \frac{\alpha_3(x)}{\alpha_2(x)}.$$

The pole (in x) of the left-hand side depends on $z \in \Theta$, whereas the pole of the right-hand side does not; a contradiction. Lemma 4 is proven.

Lemma 5. $\mathcal{D}(\psi, \psi_{c'})$ is a free $\mathcal{D}$ -module of rank 2.

PROOF. Suppose that $\mathcal{D}(\psi, \psi_{c'})$ is not a free $\mathcal{D}$ -module. Then there exist operators $d_1, d_2 \in \mathcal{D}$ such that

$$d_1 \psi_{c'} + d_2 \psi = 0. \quad (3)$$

Consider the case in which $\text{ord}(d_1) > \text{ord}(d_2) - 1$, where ord is the order of the operator. Suppose that $\text{ord}(d_1) = n$. The operator d_1 looks like

$$d_1 = f_n(x) \partial_{x_1}^n + f_{n-1}(x) \partial_{x_1}^{n-1} \partial_{x_2} + \dots + f_0(x) \partial_{x_2}^n + \dots,$$

where $f_j(x) \in \mathcal{O}$, $j = 0, \dots, n$. Divide (3) by

$$\exp(-x_1 \partial_{z_1} \log \theta(z) - x_2 \partial_{z_2} \log \theta(z)),$$

multiply by $\theta^{n+2}(z)$, and assume $z \in \Theta$. We obtain

$$\theta(z+c+c'+x)\theta(z-c')(f_n(x)\theta_1^n(z)+f_{n-1}(x)\theta_1^{n-1}(z)\theta_2(z)+\dots+f_0(x)\theta_2^n(z)) = 0. \quad (4)$$

By Lemma 3, there is a point $p \in \mathbf{C}^2$ such that $p \in \Theta$ and $\theta_1(p) = 0$. Put $z = p$ in (4). We obtain $f_0 = 0$. Divide (4) by $\theta_1(z)$ and put again $z = p$. We obtain $f_1 = 0$. Proceeding similarly, we obtain $f_n = f_{n-1} = \dots =$

$f_0 = 0$. Consequently, the inequality $\text{ord}(d_1) > \text{ord}(d_2) - 1$ is impossible. Similarly, we show that the inequality $\text{ord}(d_1) + 1 < \text{ord}(d_2)$ is impossible either. Consider the case in which $\text{ord}(d_1) + 1 = \text{ord}(d_2) = n$. Let the operators d_1 and d_2 look like

$$d_1 = f_{n-1}(x)\partial_{x_1}^{n-1} + f_{n-2}(x)\partial_{x_1}^{n-2}\partial_{x_2} + \dots + f_0(x)\partial_{x_2}^{n-1} + \dots,$$

$$d_2 = g_n(x)\partial_{x_1}^n + g_{n-1}(x)\partial_{x_1}^{n-1}\partial_{x_2} + \dots + g_0(x)\partial_{x_2}^n + \dots$$

Divide (3) by

$$\exp(-x_1\partial_{z_1}\log\theta(z) - x_2\partial_{z_2}\log\theta(z)),$$

multiply by $\theta^{n+1}(z)$, and assume $z \in \Theta$. Then

$$\theta(z+c+c'+x)\theta(z-c')(f_{n-1}(x)\theta_1^{n-1}(z)+f_{n-2}(x)\theta_1^{n-2}(z)\theta_2(z)+\dots+f_0(x)\theta_2^{n-1}(z))$$

$$-\theta(z+c+x)(g_n(x)\theta_1^n(z)+g_{n-1}(x)\theta_1^{n-1}(z)\theta_2(z)+\dots+g_0(x)\theta_2^n(z))=0.$$

Decompose

$$f_{n-1}(x)\theta_1^{n-1}(z)+f_{n-2}(x)\theta_1^{n-2}(z)\theta_2(z)+\dots+f_0(x)\theta_2^{n-1}(z),$$

$$g_n(x)\theta_1^n(z)+g_{n-1}(x)\theta_1^{n-1}(z)\theta_2(z)+\dots+g_0(x)\theta_2^n(z)$$

into factors to obtain

$$\theta(z+c+c'+x)\theta(z-c')(a_{n-1}(x)\theta_1(z)+b_{n-1}(x)\theta_2(z))\dots(a_1(x)\theta_1(z)+b_1(x)\theta_2(z))$$

$$-\theta(z+c+x)(a'_n(x)\theta_1(z)+b'_n(x)\theta_2(z))\dots(a'_1(x)\theta_1(z)+b'_1(x)\theta_2(z))=0, \quad (5)$$

where $a'_j(x)$, $b'_j(x)$, $a_k(x)$, and $b_k(x)$ are some functions, $j = 1, \dots, n$, $k = 1, \dots, n-1$.

Fix a point x . Observe that if two functions $a'_j\theta_1(z)+b'_j\theta_2(z)$ and $a_k\theta_1(z)+b_k\theta_2(z)$ have a common zero on Θ then they are proportional. Indeed, suppose the contrary. Then the ratio of these functions has a single simple pole on $\Theta \subset X^2$ (by Lemma 3). Consequently, the mapping

$$(a'_j\theta_1(z)+b'_j\theta_2(z):a_k\theta_1(z)+b_k\theta_2(z)):\Theta\rightarrow\mathbf{CP}^1,$$

where $\mathbf{CP}^1$ is the one-dimensional projective space, is an isomorphism. Since the genus of the Riemann surface $\Theta \subset X^2$ equals 2, this is a contradiction.

The zeros of $a_k\theta_1(z) + b_k\theta_2(z)$ and $\theta(z+c+x)$ on Θ cannot coincide, since these functions can be regarded as sections of the line bundle over $\Theta \subset X^2$ and their ratio

$$\frac{a_k\theta_1(z) + b_k\theta_2(z)}{\theta(z+c+x)} \quad (6)$$

is again a section of the line bundle but it has neither poles nor zeros. The only sections with such properties are constants, but (6) is not constant on Θ .

Consequently, (5) amounts to the equality

$$\theta(z+c+c'+x)\theta(z-c') + \theta(z+c+x)(a(x)\theta_1(z) + b(x)\theta_2(z)) = 0,$$

where $a(x)$ and $b(x)$ are some functions. We conclude that the functions $\theta(z-c')$ and $\theta(z+c+x)$ must have a common zero on Θ for every x , which is impossible as we easily see, for instance, from Riemann's theorem about the zeros of a theta function on a Riemann surface. Lemma 5 is proven.

To complete the proof of Nakayashiki's theorem, it suffices to demonstrate that the $\mathcal{D}$ -modules $\mathcal{D}(\psi, \psi_{c'})$ and M_c coincide.

Lemma 6. *The following equality holds: $\mathcal{D}(\psi, \psi_{c'}) = M_c$.*

PROOF. The inclusion $\mathcal{D}(\psi, \psi_{c'}) \subset M_c$ is obvious. Fix a point x . Denote by $\mathcal{D}^k(\psi, \psi_{c'}) \subset \mathcal{D}(\psi, \psi_{c'})$ the set of functions

$$\{d_1\psi_{c'} + d_2\psi \mid d_1, d_2 \in \mathcal{D}, \text{ord}(d_1) \leq k-2, \text{ord}(d_2) \leq k-1\}.$$

To prove the lemma, it suffices to demonstrate that

$$\dim_{\mathbf{C}} M_c(k) = \dim_{\mathbf{C}} \mathcal{D}^k(\psi, \psi_{c'}).$$

By Lemma 1, we have

$$\dim_{\mathbf{C}} M_c(k) = \dim_{\mathbf{C}} \left\{ \sum_{a \in \mathbf{Z}^g / k\mathbf{Z}^g} \beta_a \theta \left[\frac{a}{k}, 0 \right] (kz + c + x, k\Omega) \mid \beta_a \in \mathbf{C} \right\}.$$

Consequently, $\dim_{\mathbf{C}} M_c(k) = k^2$ (see, for instance, [9]). Since the dimension of the space of differential operators of order $\leq k$ with constant coefficients equals

$$1 + \dots + (k+1) = \frac{(k+1)(k+2)}{2}$$

and the $\mathcal{D}$ -module $\mathcal{D}(\psi, \psi_{c'})$ is free, we have

$$\dim_{\mathbf{C}} \mathcal{D}^k(\psi, \psi_{c'}) = \frac{(k-1)k}{2} + \frac{k(k+1)}{2} = k^2.$$

Lemma 6 is proven.

4 A Commutative Ring of (2×2) -Matrix Differential Operators

Denote by D the ring of differential operators $\mathbf{C}[\partial_{z_1}, \dots, \partial_{z_g}]$ with constant coefficients. By definition, M_c is a D -module. Let $\Phi_c = (\phi_{1c}, \dots, \phi_{g!c})^\top$ be a basis for M_c . Then we have the ring embedding

$$\mathcal{L}_{\Phi_c} : D \rightarrow \text{Mat}(g!, \mathcal{D})$$

defined by the formula

$$\mathcal{L}_{\Phi_c}(d)\Phi_c = d\Phi_c,$$

where $d \in D$ and $d\Phi_c = (d\phi_{1c}, \dots, d\phi_{g!c})^\top$. The range of $\mathcal{L}_{\Phi_c}$ is a commutative ring of matrix differential operators which is isomorphic to D . Denote the operator $\mathcal{L}_{\Phi_c}(\partial_{z_j})$ by Z_j , $j = 1, \dots, g$.

Find the commutator $[L_{\Phi_c}(\lambda), Z_j]$ for $\lambda \in A_\Theta$:

$$Z_j L_{\Phi_c}(\lambda)\Phi_c = Z_j(\lambda\Phi_c) = \lambda(Z_j\Phi_c) = \lambda(\partial_{z_j}\Phi_c),$$

$$L_{\Phi_c}(\lambda)Z_j\Phi_c = L_{\Phi_c}\partial_{z_j}\Phi_c = \partial_{z_j}(L_{\Phi_c}(\lambda)\Phi_c) = \partial_{z_j}(\lambda\Phi_c) = (\partial_{z_j}\lambda)\Phi_c + \lambda(\partial_{z_j}\Phi_c).$$

Thus,

$$[L_{\Phi_c}(\lambda), Z_j]\Phi_c = (\partial_{z_j}\lambda)\Phi_c,$$

which proves the following

Proposition 1. $[L_{\Phi_c}(\lambda), Z_j] = L_{\Phi_c}(\partial_{z_j}\lambda)$ for $\lambda \in A_\Theta$, $j = 1, \dots, g$.

We now clarify how the matrix operators change when we replace a basis for the $\mathcal{D}$ -module M_c . Suppose that $\Phi_c = (\phi_{1c}, \dots, \phi_{g!c})^\top$ and $\Psi_c = (\psi_{1c}, \dots, \psi_{g!c})^\top$ are two bases for the $\mathcal{D}$ -module M_c . By Nakayashiki's theorem, there exist unique operators $\mathcal{A}, \mathcal{B} \in \text{Mat}(g!, \mathcal{D})$ such that $\mathcal{A}\Phi_c = \Psi_c$ and $\mathcal{B}\Psi_c = \Phi_c$; hence, $\mathcal{A}\mathcal{B} = \mathcal{B}\mathcal{A}$ is the identity in $\text{Mat}(g!, \mathcal{D})$. We denote the operator $\mathcal{B}$ by $\mathcal{A}^{-1}$. It is easy to verify that

$$\mathcal{A}L_{\Phi_c}(\lambda)\mathcal{A}^{-1}\Psi_c = \lambda\Psi_c, \quad \mathcal{A}\mathcal{L}_{\Phi_c}(d)\mathcal{A}^{-1}\Psi_c = d\Psi_c,$$

where $\lambda \in A_\Theta$ and $d \in D$. Consequently, we have the following

Lemma 7 *The equalities hold:*

$$\mathcal{A}L_{\Phi_c}(\lambda)\mathcal{A}^{-1} = L_{\Psi_c}(\lambda), \quad \mathcal{A}\mathcal{L}_{\Phi_c}(d)\mathcal{A}^{-1} = \mathcal{L}_{\Psi_c}(d).$$

Henceforth we suppose that $g = 2$. In Proposition 2 we find the second-order operators $L_{\Phi_c}(\lambda)$. They correspond to the following spectral functions:

$$\lambda = \frac{\partial^2 \log \theta(z)}{\partial z_k \partial z_j}, \quad j, k = 1, 2.$$

In Proposition 3 we find the operators Z_1 and Z_2 . In Lemma 9 we prove that the operators of Propositions 1–3 generate the ring $L_{\Phi_c}(A_\Theta)$.

The following lemma can be proven by straightforward calculations which we omit:

Lemma 8. *The equality holds:*

$$\begin{aligned} & \partial_{x_k} \partial_{x_j} \left(\frac{\theta(z + c + x)}{\theta(z)} \exp(-x_1 \partial_{z_1} \log \theta(z) - x_2 \partial_{z_2} \log \theta(z)) \right) \\ &= \partial_{z_k} \partial_{z_j} \left(\frac{\theta(z + c + x)}{\theta(z)} \right) \exp(-x_1 \partial_{z_1} \log \theta(z) - x_2 \partial_{z_2} \log \theta(z)) \\ &+ \partial_{z_k} \partial_{z_j} (\log \theta(z)) \frac{\theta(z + c + x)}{\theta(z)} \exp(-x_1 \partial_{z_1} \log \theta(z) - x_2 \partial_{z_2} \log \theta(z)), \end{aligned}$$

where $k, j = 1, 2$.

Let c' be in a general position. Denote by $p_1, p_2 \in \mathbf{C}^2$ the intersection points of $\Gamma_{c'}$ with Θ (p_1 and p_2 are defined to within elements of the lattice $\mathbf{Z}^2 + \Omega \mathbf{Z}^2$). Denote by $L_{c,c'}$ the Nakayashiki operators for the basis $\psi, \psi_{c'}$. For brevity, we introduce the notations

$$H_{c,c'}^{kj} = [L_{c,c'}(\partial_{z_k} \partial_{z_j} \log \theta(z))]_{11}, \quad F_{c,c'}^{kj} = [L_{c,c'}(\partial_{z_k} \partial_{z_j} \log \theta(z))]_{12}.$$

Proposition 2. *The equality holds:*

$$[L_{c,c'}(\partial_{z_k} \partial_{z_j} \log \theta(z))]_{11} = -\partial_{x_k} \partial_{x_j} + f_{c,c'}^{kj} \partial_{x_1} + g_{c,c'}^{kj} \partial_{x_2} + h_{c,c'}^{kj},$$

with

$$\begin{aligned} f_{c,c'}^{kj} &= \frac{\theta_2(p_1) \theta_2(p_2)}{\theta_2(p_2) \theta_1(p_1) - \theta_2(p_1) \theta_1(p_2)} \\ &\times \left(\partial_{x_k} \log \theta(p_1 + c + x) \frac{\theta_j(p_1)}{\theta_2(p_1)} + \partial_{x_j} \log \theta(p_1 + c + x) \frac{\theta_k(p_1)}{\theta_2(p_1)} - \frac{\theta_{kj}(p_1)}{\theta_2(p_1)} \right. \\ &\left. - \partial_{x_k} \log \theta(p_2 + c + x) \frac{\theta_j(p_2)}{\theta_2(p_2)} - \partial_{x_j} \log \theta(p_2 + c + x) \frac{\theta_k(p_2)}{\theta_2(p_2)} + \frac{\theta_{kj}(p_2)}{\theta_2(p_2)} \right), \end{aligned}$$

$$g_{c,c'}^{kj} = \frac{\theta_1(p_1)\theta_1(p_2)}{\theta_2(p_1)\theta_1(p_2) - \theta_2(p_2)\theta_1(p_1)} \\ \times \left(\partial_{x_j} \log \theta(p_1 + c + x) \frac{\theta_k(p_1)}{\theta_1(p_1)} + \partial_{x_k} \log \theta(p_1 + c + x) \frac{\theta_j(p_1)}{\theta_1(p_1)} - \frac{\theta_{kj}(p_1)}{\theta_1(p_1)} \right. \\ \left. - \partial_{x_j} \log \theta(p_2 + c + x) \frac{\theta_k(p_2)}{\theta_1(p_2)} - \partial_{x_k} \log \theta(p_2 + c + x) \frac{\theta_j(p_2)}{\theta_1(p_2)} + \frac{\theta_{kj}(p_2)}{\theta_1(p_2)} \right),$$

$$h_{c,c'}^{kj} = \frac{\theta(\Delta + c')}{\theta(\Delta + c' + c + x)} (\partial_{z_k} \partial_{z_j} - f_{c,c'}^{kj} \partial_{z_1} - g_{c,c'}^{kj} \partial_{z_2} + 2 \partial_{z_k} \partial_{z_j} \log \theta(z)) \\ \times \left(\frac{\theta(z + c + x)}{\theta(z)} \right) \Big|_{z=\Delta+c'},$$

$$[L_{c,c'}(\partial_{z_k} \partial_{z_j} \log \theta(z))]_{12} = \frac{\theta^2(0)}{\theta(c + c' + x)\theta(c')} \\ \times (\partial_{z_k} \partial_{z_j} - f_{c,c'}^{kj} \partial_{z_1} - g_{c,c'}^{kj} \partial_{z_2} - h_{c,c'}^{kj} + 2 \partial_{z_k} \partial_{z_j} \log \theta(z)) \left(\frac{\theta(z + c + x)}{\theta(z)} \right) \Big|_{z=0},$$

$$[L_{c,c'}(\partial_{z_k} \partial_{z_j} \log \theta(z))]_{21} = F_{c+c',-c'}^{kj} (\alpha_{11} H_{c,c'}^{11} + \alpha_{12} H_{c,c'}^{12} + \alpha_{22} H_{c,c'}^{22} + \alpha),$$

$$[L_{c,c'}(\partial_{z_k} \partial_{z_j} \log \theta(z))]_{22} = F_{c+c',-c'}^{kj} (\alpha_{11} F_{c,c'}^{11} + \alpha_{12} F_{c,c'}^{12} + \alpha_{22} F_{c,c'}^{22}) + H_{c+c',-c'}^{kj},$$

where $\alpha, \alpha_{kj} \in \mathbf{C}$ are such that

$$\frac{\theta(z - c')\theta(z + c')}{\theta^2(z)} = \alpha_{11} \partial_{z_1}^2 \log \theta(z) + \alpha_{12} \partial_{z_1} \partial_{z_2} \log \theta(z) + \alpha_{22} \partial_{z_2}^2 \log \theta(z) + \alpha.$$

PROOF OF PROPOSITION 2. It is easy to verify that the function

$$\frac{\partial_{x_k} \partial_{x_j} \psi + \partial_{z_k} \partial_{z_j} \log \theta(z) \psi}{\exp(-x_1 \partial_{z_1} \log \theta(z) - x_2 \partial_{z_2} \log \theta(z))}$$

has a second-order pole on Θ ; i.e.,

$$\partial_{x_k} \partial_{x_j} \psi + \partial_{z_k} \partial_{z_j} \log \theta(z) \psi \in M_c(2).$$

Hence, the operator $H_{c,c'}^{kj}$ looks like

$$H_{c,c'}^{kj} = -\partial_{x_k} \partial_{x_j} + f_{c,c'}^{kj} \partial_{x_1} + g_{c,c'}^{kj} \partial_{x_2} + h_{c,c'}^{kj},$$

and $F_{c,c'}^{kj}$ is the operator of multiplication by the function

$$H_{c,c'}^{kj} \psi + F_{c,c'}^{kj} \psi_{c'} = \partial_{z_k} \partial_{z_j} \log \theta(z) \psi. \quad (7)$$

Rewrite (7) for $z \in \Gamma_{c'}$ as follows:

$$H_{c,c'}^{kj} \psi = \partial_{z_k} \partial_{z_j} \log \theta(z) \psi.$$

By Lemmas 2 and 8, the last equality amounts to

$$(\partial_{z_k} \partial_{z_j} - f_{c,c'}^{kj} \partial_{z_1} - g_{c,c'}^{kj} \partial_{z_2} - h_{c,c'}^{kj} + 2\partial_{z_k} \partial_{z_j} \log \theta(z)) \left(\frac{\theta(z+c+x)}{\theta(z)} \right) = 0. \quad (8)$$

Multiply both sides of (8) by $\theta^2(z)$ and put $z = p_1$ and $z = p_2$ to obtain the following system of linear equations in $f_{c,c'}^{kj}$ and $g_{c,c'}^{kj}$:

$$\begin{aligned} f_{c,c'}^{kj} \theta_1(p_1) + g_{c,c'}^{kj} \theta_2(p_1) &= \frac{\theta_j(p_1+c+x) \theta_k(p_1) + \theta_k(p_1+c+x) \theta_j(p_1) - \theta_{kj}(p_1)}{\theta(p_1+c+x)}, \\ f_{c,c'}^{kj} \theta_1(p_2) + g_{c,c'}^{kj} \theta_2(p_2) &= \frac{\theta_j(p_2+c+x) \theta_k(p_2) + \theta_k(p_2+c+x) \theta_j(p_2) - \theta_{kj}(p_2)}{\theta(p_2+c+x)}. \end{aligned}$$

Solving it, we find $f_{c,c'}^{kj}$ and $g_{c,c'}^{kj}$. Putting $z = \Delta + c'$ in (8) ($\Delta + c' \in \Gamma_{c'}$, since $\theta(\Delta) = 0$), we find $h_{c,c'}^{kj}$. It follows from Lemmas 2 and 8 that (7) is equivalent to the equality

$$\begin{aligned} &(-\partial_{z_k} \partial_{z_j} + f_{c,c'}^{kj} \partial_{z_1} + g_{c,c'}^{kj} \partial_{z_2} + h_{c,c'}^{kj}) \left(\frac{\theta(z+c+x)}{\theta(z)} \right) \\ &+ F_{c,c'}^{kj} \frac{\theta(z+c+c'+x) \theta(z-c')}{\theta^2(z)} = 2\partial_{z_i} \partial_{z_j} \log \theta(z) \frac{\theta(z+c+x)}{\theta(z)}. \end{aligned}$$

Putting $z = 0$ in this equality, we obtain $F_{c,c'}^{kj}$.

We find the remaining entries of the operator $L_{c,c'}(\partial_{z_k} \partial_{z_j} \log \theta(z))$. Replace c' with $-c'$ and c with $c+c'$ in (7) and multiply both sides by $\frac{\theta(z-c')}{\theta(z)}$ to obtain

$$H_{c+c',-c'}^{kj} \psi_{c'} + F_{c+c',-c'}^{kj} \frac{\theta(z-c') \theta(z+c')}{\theta^2(z)} \psi = \partial_{z_k} \partial_{z_j} \log \theta(z) \psi_{c'}. \quad (9)$$

Let

$$\frac{\theta(z - c')\theta(z + c')}{\theta^2(z)} = \alpha_{11}\partial_{z_1}^2 \log \theta(z) + \alpha_{12}\partial_{z_1}\partial_{z_2} \log \theta(z) + \alpha_{22}\partial_{z_2}^2 \log \theta(z) + \alpha,$$

where $\alpha, \alpha_{kj} \in \mathbf{C}$. The existence of the constants α, α_{kj} , $k, j = 1, 2$, follows from the fact that the dimension of the space of meromorphic functions with a pole of order at most 2 on Θ equals 4 (see, for instance, [9]). The functions 1 and $\partial_{z_k}\partial_{z_j} \log \theta(z)$, $k, j = 1, 2$, are linearly independent over $\mathbf{C}$, which can be shown, for instance, by means of Nakayashiki's theorem. The operators $L_{c,c'}(1)$ and $L_{c,c'}(\partial_{z_k}\partial_{z_j} \log \theta(z))$, $k, j = 1, 2$, are linearly independent, since the leading part of the 11-entries of these operators, as demonstrated, are equal respectively to 1 and $-\partial_{x_k}\partial_{x_j}$. Now, (9) amounts to

$$F_{c+c',-c'}^{kj}(\alpha_{11}H_{c,c'}^{11} + \alpha_{12}H_{c,c'}^{12} + \alpha_{22}H_{c,c'}^{22} + \alpha)\psi$$

$$+ (F_{c+c',-c'}^{kj}(\alpha_{11}F_{c,c'}^{11} + \alpha_{12}F_{c,c'}^{12} + \alpha_{22}F_{c,c'}^{22}) + H_{c+c',-c'}^{kj})\psi_{c'} = \partial_{z_k}\partial_{z_j} \log \theta(z)\psi_{c'};$$

whence we obtain $[L_{c,c'}(\partial_{z_k}\partial_{z_j} \log \theta(z))]_{21}$ and $[L_{c,c'}(\partial_{z_k}\partial_{z_j} \log \theta(z))]_{22}$, completing the proof of Proposition 2.

We find the operators Z_1 and Z_2 .

Proposition 3. *The operators Z_1 and Z_2 for the basis $\psi, \psi_{c'}$ look like*

$$[Z_j]_{11} = -x_1H_{c,c'}^{1j} - x_2H_{c,c'}^{j2} + \partial_{x_j}, \quad [Z_j]_{12} = -x_1F_{c,c'}^{1j} - x_2F_{c,c'}^{j2},$$

$$[Z_j]_{21} = -x_1[L_{c,c'}(\partial_{z_1}\partial_{z_j} \log \theta(z))]_{21}$$

$$-x_2[L_{c,c'}(\partial_{z_j}\partial_{z_2} \log \theta(z))]_{21} + k_1^j\partial_{x_1} + k_2^j\partial_{x_2} + h_{c,c'}^j,$$

$$[Z_j]_{22} = -x_1[L_{c,c'}(\partial_{z_1}\partial_{z_j} \log \theta(z))]_{22} - x_2[L_{c,c'}(\partial_{z_j}\partial_{z_2} \log \theta(z))]_{22} + g_{c,c'}^j + 2\partial_{x_j},$$

where

$$\begin{aligned} k_1^j &= \frac{1}{\theta_1(p_2)\theta_2(p_1) - \theta_1(p_1)\theta_2(p_2)} \\ &\times \left(\frac{\theta_j(p_1 - c')\theta(p_1 + c + x + c')\theta_2(p_2)}{\theta(p_1 + c + x)} - \frac{\theta_j(p_2 - c')\theta(p_2 + c + x + c')\theta_2(p_1)}{\theta(p_2 + c + x)} \right), \\ k_2^j &= \frac{1}{\theta_1(p_1)\theta_2(p_2) - \theta_1(p_2)\theta_2(p_1)} \\ &\times \left(\frac{\theta_j(p_1 - c')\theta(p_1 + c + x + c')\theta_1(p_2)}{\theta(p_1 + c + x)} - \frac{\theta_j(p_2 - c')\theta(p_2 + c + x + c')\theta_1(p_1)}{\theta(p_2 + c + x)} \right), \end{aligned}$$

$$\begin{aligned}
h_{c,c'}^j(x) &= \frac{\theta_j(\Delta)\theta(\Delta+c+x+2c')}{\theta(\Delta+c')\theta(\Delta+c+x+c')} \\
&-k_1^j \left(\frac{\theta_1(\Delta+c+x+c')}{\theta(\Delta+c+x+c')} - \frac{\theta_1(\Delta+c')}{\theta(\Delta+c')} \right) - k_2^j \left(\frac{\theta_2(\Delta+c'+c+x)}{\theta(\Delta+c'+c+x)} - \frac{\theta_2(\Delta+c')}{\theta(\Delta+c')} \right), \\
g_{c,c'}^j(x) &= -\frac{\theta_j(c')}{\theta(c')} + \frac{\theta_j(c+x+c')}{\theta(c+x+c')} \\
&- \frac{\theta^2(0)}{\theta(c')\theta(c+x+c')} (k_1^j \partial_{z_1} + k_2^j \partial_{z_2} + h_{c,c'}^j) \left(\frac{\theta(z+c+x)}{\theta(z)} \right) \Big|_{z=0},
\end{aligned}$$

$j = 1, 2$.

PROOF OF PROPOSITION 3 We have to find an operator $Z_j \in \text{Mat}(2, \mathcal{D})$ such that

$$\begin{aligned}
[Z_j]_{11}\psi + [Z_j]_{12}\psi_{c'} &= \partial_{z_j}\psi, \quad [Z_j]_{21}\psi + [Z_j]_{22}\psi_{c'} = \partial_{z_j}\psi_{c'}, \\
\partial_{z_j}\psi &= \partial_{z_j} \left(\frac{\theta(z+c+x)}{\theta(z)} \right) \exp(-x_1 \partial_{z_1} \log \theta(z) - x_2 \partial_{z_2} \log \theta(z)) \\
&- (x_1 \partial_{z_1} \partial_{z_j} \log \theta(z) + x_2 \partial_{z_j} \partial_{z_2} \log \theta(z)) \frac{\theta(z+c+x)}{\theta(z)} \exp(-x_1 \partial_{z_1} \log \theta(z) \\
&- x_2 \partial_{z_2} \log \theta(z)) = (-x_1 H_{c,c'}^{1j} - x_2 H_{c,c'}^{j2} + \partial_{x_j})\psi - (x_1 F_{c,c'}^{1j} + x_2 F_{c,c'}^{j2})\psi_{c'}.
\end{aligned}$$

Therefore,

$$[Z_j]_{11} = -x_1 H_{c,c'}^{1j} - x_2 H_{c,c'}^{j2} + \partial_{x_j}, \quad [Z_j]_{12} = -x_1 F_{c,c'}^{1j} - x_2 F_{c,c'}^{j2}.$$

We find the remaining entries of Z_j :

$$\begin{aligned}
\partial_{z_j}\psi_{c'} &= \partial_{z_j} \left(\frac{\theta(z-c')}{\theta(z)} \right) \frac{\theta(z+c+x+c')}{\theta(z)} \exp(-x_1 \partial_{z_1} \log \theta(z) - x_2 \partial_{z_2} \log \theta(z)) \\
&+ \frac{\theta(z-c')}{\theta(z)} \partial_{z_j} \left(\frac{\theta(z+c+x+c')}{\theta(z)} \right) \exp(-x_1 \partial_{z_1} \log \theta(z) - x_2 \partial_{z_2} \log \theta(z)) \\
&- (x_1 \partial_{z_1} \partial_{z_j} \log \theta(z) + x_2 \partial_{z_j} \partial_{z_2} \log \theta(z)) \frac{\theta(z-c')\theta(z+c+x+c')}{\theta^2(z)} \\
&\times \exp(-x_1 \partial_{z_1} \log \theta(z) - x_2 \partial_{z_2} \log \theta(z)). \tag{10}
\end{aligned}$$

Immediate calculations show that

$$\begin{aligned} \partial_{z_j} \left(\frac{\theta(z - c')}{\theta(z)} \right) \frac{\theta(z + c + x + c')}{\theta(z)} \exp(-x_1 \partial_{z_1} \log \theta(z) - x_2 \partial_{z_2} \log \theta(z)) - \partial_{x_j} \psi_{c'} \\ = \frac{\theta_j(z - c') \theta(z + c + x + c') - \theta(z - c') \theta_j(z + c + x + c')}{\theta^2(z)} \\ \times \exp(-x_1 \partial_{z_1} \log \theta(z) - x_2 \partial_{z_2} \log \theta(z)) \in M_c(2); \end{aligned}$$

consequently, there exist functions $k_1^j, k_2^j, h_{c,c'}^j, g_{c,c'}^j \in \mathcal{O}$ such that

$$\begin{aligned} \partial_{z_j} \left(\frac{\theta(z - c')}{\theta(z)} \right) \frac{\theta(z + c + x + c')}{\theta(z)} \exp(-x_1 \partial_{z_1} \log \theta(z) - x_2 \partial_{z_2} \log \theta(z)) \\ = (k_1^j \partial_{x_1} + k_2^j \partial_{x_2} + h_{c,c'}^j) \psi + (g_{c,c'}^j + \partial_{x_j}) \psi_{c'}. \end{aligned} \quad (11)$$

To simplify notation, we omit the subscripts c and c' of k_1^j and k_2^j . We multiply both sides of (11) by $\theta^2(z)$ and put $z = p_1$ and $z = p_2$. We obtain the following system of linear equations in k_1^j and k_2^j :

$$\begin{aligned} k_1^j \theta_1(p_1) + k_2^j \theta_2(p_1) &= - \frac{\theta_j(p_1 - c') \theta(p_1 + c + x + c')}{\theta(p_1 + c + x)}, \\ k_1^j \theta_1(p_2) + k_2^j \theta_2(p_2) &= - \frac{\theta_j(p_2 - c') \theta(p_2 + c + x + c')}{\theta(p_2 + c + x)}. \end{aligned}$$

Solving it, we find k_1^j and k_2^j . Putting $z = \Delta + c'$ in (11) ($\Delta + c' \in \Gamma_{c'}$), we find $h_{c,c'}^j(x)$. Putting $z = 0$ in (11), we find $g_{c,c'}^j(x)$. From (10) and (11) we obtain $[Z_j]_{21}$ and $[Z_j]_{22}$. Proposition 3 is proven.

Let us show that the so-obtained operators generate the ring $L_{c,c'}(A_\Theta)$.

Lemma 9. *The operators*

$$L_{\Phi_c}(\partial_{z_k} \partial_{z_j} \log \theta(z)), \quad L_{\Phi_c}(\partial_{z_k} \partial_{z_j} \partial_{z_s} \log \theta(z)),$$

where $s, j, k = 1, 2$, generate the operator ring $L_{c,c'}(A_\Theta)$.

PROOF. The dimension of the space of meromorphic functions on X^2 with pole on Θ of multiplicity at most 3 equals 9 (see, for instance, [9]). Take a basis for this space which consists of functions of the form

$$1, \frac{f_1}{\theta^3}, \dots, \frac{f_8}{\theta^3},$$

where f_i is not divisible by θ , $i = 1, \dots, 8$. As we show below, we can take the basis of the functions

$$\begin{aligned} 1, \quad \partial_{z_k} \partial_{z_j} \partial_{z_s} \log \theta(z), \quad \partial_{z_1}^3 \log \theta(z) + \partial_{z_j} \partial_{z_s} \log \theta(z), \\ (\partial_{z_1} \partial_{z_2} \log \theta(z))^2 - \partial_{z_1}^2 \log \theta(z) \partial_{z_2}^2 \log \theta(z), \end{aligned} \quad (12)$$

where $k, j, s = 1, 2$. By the Lefschetz theorem, the mapping $(\theta^3 : f_1 : \dots : f_8)$ defines an embedding $F : X^2 \rightarrow \mathbf{CP}^8$ into the projective space. Let $(y_0 : \dots : y_8)$ be homogeneous coordinates in $\mathbf{CP}^8$. Then the restrictions of the functions $\frac{y_1}{y_0}, \dots, \frac{y_8}{y_0}$ to $F(X^2)$ generate the coordinate ring of the affine algebraic variety $F(X^2 \setminus \Theta)$. Consequently, the functions

$$\frac{f_1}{\theta^3}, \dots, \frac{f_8}{\theta^3}$$

generate A_Θ . We are left with demonstrating that the functions (12) are linearly independent over $\mathbf{C}$. This follows from the fact that the operators

$$\begin{aligned} L_{c,c'}(1), \quad L_{c,c'}(\partial_{z_k} \partial_{z_j} \partial_{z_s} \log \theta(z)), \quad L_{c,c'}(\partial_{z_1}^3 \log \theta(z) + \partial_{z_j} \partial_{z_s} \log \theta(z)), \\ (L_{c,c'}(\partial_{z_1} \partial_{z_2} \log \theta(z))^2 - \partial_{z_1}^2 \log \theta(z) \partial_{z_2}^2 \log \theta(z)), \end{aligned}$$

where $k, j, s = 1, 2$, are linearly independent, because the leading symbols of the 11-entries of these operators are equal respectively to

$$\begin{aligned} 1, \quad -2\partial_{x_k} \partial_{x_j} \partial_{x_s}, \quad -2\partial_{x_1}^3 - \partial_{x_j} \partial_{x_s}, \\ f_{c,c'}^{22} \partial_{x_1}^3 + (g_{c,c'}^{22} - 2f_{c,c'}^{12}) \partial_{x_1}^2 \partial_{x_2} + (f_{c,c'}^{11} - 2g_{c,c'}^{12}) \partial_{x_1} \partial_{x_2}^2 + g_{c,c'}^{11} \partial_{x_2}^3. \end{aligned}$$

Lemma 9 is proven.

References.

- [1] Nakayashiki A., Structure of Baker–Akhiezer modules of principally polarized Abelian varieties, commuting partial differential operators and associated integrable systems, *Duke Math. J.*, 1991, vol. 62, N. 2, 315–358.
- [2] Dubrovin B. A., Krichever I. M., and Novikov S. P., The Schroedinger equation in a periodic magnetic field and Riemann surfaces, *Soviet Math. Dokl.*, 1976, vol. 17, 947–952.
- [3] Veselov A. P. and Novikov S. P., Finite-gap two-dimensional Schrödinger operators. Potential operators, *Soviet Math. Dokl.*, 1984, vol. 30, 705–708.
- [4] Feldman J., Knorrer H., and Trubowitz E., There is no two-dimensional analogue of Lamé’s equation, *Math. Ann.*, 1992, vol. 294, N. 2, 295–324.

- [5] Nakayashiki A., Commuting partial differential operators and vector bundles over Abelian varieties, Amer. J. Math., 1994, vol. 116, 65–100.
- [6] Mukai S., Duality between $D(X)$ and $D(\widehat{X})$ with its application to Picard sheaves, Nagoya Math. J., 1981, vol. 81, 153–175.
- [7] Andreotti A. and Mayer A., On period relations for Abelian integrals on algebraic curves, Ann. Scuola Norm. Sup. Pisa Cl. Sci (4), 1967, vol. 21, 189–238.
- [8] Krichever I. M., The methods of algebraical geometry in the theory of nonlinear equations, Russian Math. Surveys (1977), vol. 32, no. 6, 185–213.
- [9] Taimanov I. A., Secants of Abelian varieties, theta functions, and soliton equations, Uspekhi Mat. Nauk, 1997, vol. 52, N. 1, 149–224,

Sobolev Institute of Mathematics, Novosibirsk
 mironov@math.nsc.ru